



Herbizidstrategien in Ackerbohnen in 2017

LLH-Versuchsstandort Bad Hersfeld - Eichhof



	BBCH 00 04.04.	BBCH 16 11.05.	BBCH 18 16.05.	BBCH 50 22.05.	SNK Test	Kontrolle: 45,12 dt/ha, rel. 100 GD 5%: 3,854
Kontrolle					AB	100
Boxer + Stomp Aqua	3,0 + 2,0				AB	103
Novitron	2,4				AB	106
Novitron; Betanal Expert*	2,4	1,2			B	95
Batanal Max Pro*; Betanal Maxx pro*		1,2	1,2		C	85
Betanal Expert*; Betanal Expert*					C	83
Clentiga*		1,5			AB	102
Pulsar 40* + Dash		0,3 + 1,0			AB	103
Proman*	3,0				AB	104
Beloukha*	50,0				AB	108
Striegel				x	AB	102
Kompostgabe 5 cm	x				A	112

* = keine Indikation in Ackerbohnen

80 90 100 110

■ Ertrag relativ in %



Herbizidstrategien in Ackerbohnen in 2017

LLH-Versuchsstandort Bad Hersfeld - Eichhof



Vorfrucht Wintergerste, Aussaat 30.03.2017, Sorte Fanfare, 45 Kö/m²,
Nmin 27.03.2017: 20 kg/ha; pH 6,1, P205 18, K20 16, Mgo 12
03.04.2017 Bandur 4,0
11.05.2017 Basagran 1,0
23.05.2017 Karate Zeon 0,075