

# Herbizide in Winterweizen gegen Windhalm in 2011/2012, LLH-Versuchsstandort Limburg

Varianten	l/kg/ha	HRAC Gräser	09-11 16.10.11	NAF 16.04.12	Wirkungsgrad in %
Bacara Forte	0,8	F1,F1,K3	X		100
Boxer + Beflex	2,5 + 0,4	N,F1	X		100
Boxer + Diflanil	2,5 + 0,25	N,F1	X		96
Trinity	2,0	C2,F1,K1	X		100
Fenikan	2,0	C2,F1	X		94
Carmina 640 + Alliance	2,0 + 0,05	C2,F1,B,F1	X		97
Corello + Dash (Termin zu früh)	2,0 + 1,0	B,K1	X		99
Sumimax + Arelon Top	0,06 + 1,5	E,C2	X		99
Sumimax + Herold SC	0,06 + 0,2	E,F1,K3	X		100
Absolute M	0,2	B	x		96
Broadway + Netzmittel	0,13 + 0,6	B		X	89
Axial Komplett	1,0	A		X	100
Axial Komplett + Arelon Top	1,0 + 1,5	A,C2		X	100
Atlantis OD + Starane XL	0,85 + 1,0	B		X	98
Ralon Super + Lentipur 700 + Mero	1,0 + 3,0	A,C2		X	100
Arelon Top + Concert SX	2,0 + 0,1	C2,B		X	73
Arelon Top + Fox + Primus	2,0 + 1,0 + 0,075	C2		x	83

Aussaat: 29.09.2011, Sorte Potential, Kontrolle 132 Windhalmrispen/m<sup>2</sup>

■ Wirkungsgrad in %